

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 1 078 926 A1**

(12)

# EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:  
28.02.2001 Bulletin 2001/09

(21) Application number: 00907971.6

(22) Date of filing: 09.03.2000

(51) Int. Cl.<sup>7</sup>: **C07D 407/06**, C07D 405/14,  
C07D 407/14, C07D 417/14,  
C07D 455/04, C07D 491/16,  
C09B 57/02, C08F 2/50,  
G03C 1/73, G03F 7/031,  
G03H 1/02

(86) International application number:  
**PCT/JP00/01437**

(87) International publication number:  
**WO 00/53598 (14.09.2000 Gazette 2000/37)**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

(30) Priority: 09.03.1999 JP 6192299  
12.07.1999 JP 19729799  
08.03.2000 JP 2000063867

(71) Applicant:  
**Kabushiki Kaisha Hayashibara Seibutsu Kagaku  
Kenkyujo  
Okayama-shi, Okayama 700-0907 (JP)**

(72) Inventors:  
• **SATSUKI, Makoto**  
Tsukubo-gun, Okayama 701-0300 (JP)  
• **SHINPO, Akira**  
Oku-gun, Okayama 701-4301 (JP)

- **OOGA, Yasuyo**  
Kurashiki-shi, Okayama 711-0927 (JP)
- **SUGA, Sadaharu**  
Tamano-shi, Okayama 706-0011 (JP)
- **ODA, Atsushi,**  
Nippon Denki Kabushiki Gaisha-nai  
Kawasaki-shi, Kanagawa 216-8555 (JP)
- **TADA, Hiroshi,**  
Nippon Denki Kabushiki Gaisha-nai  
Kawasaki-shi, Kanagawa 216-8555 (JP)
- **SAKAGUCHI, Yoshikazu,**  
Nippon Denki K. Gaisha-nai  
Kawasaki-shi, Kanagawa 216-8555 (JP)

(74) Representative:  
**Daniels, Jeffrey Nicholas et al**  
Page White & Farrer  
54 Doughty Street  
London WC1N 2LS (GB)

## (54) PYRAN DERIVATIVES

(57) Disclosed are pyran derivatives, which have distinctive sensitivity to visible light and distinctive luminous ability. The derivatives are useful in photopolymerization, electroluminescence, and dye lasers, and are obtainable through a step of reacting a compound having a 4-cyanomethylene-2-methyl-4H-pyran skeleton with a compound having a 3-formylcoumarin skeleton.

BEST AVAILABLE COPY

EP 1 078 926 A1

**Description**TECHNICAL FIELD

- 5 [0001] The present invention relates to functional organic compounds, and more particularly, to novel pyran derivatives which are useful in photochemical polymerization, dye laser, and electroluminescence.

BACKGROUND ART

- 10 [0002] After coming into this information age, photochemical polymerization has been frequently used in a variety of fields, and beyond the bounds of the field of synthetic resins, it is being extensively used in other fields of information recordings and electric equipments such as paints, plates for printings, printing circuits, integrated circuits, etc. Photochemical polymerization, a technique for polymerizing polymerizable compounds by irradiating light, is classified roughly into photopolymerization where polymerizable compounds are initiated to be polymerized and activated by  
15 directly irradiating light, and photosensitizing polymerization where polymerizable compounds are polymerized by irradiating light in the presence of photosensitizers to form growth active-cores of photosensitizers. The photochemical polymerizations have characteristics that the initiation and suspension of polymerization can be controlled by flashing excitation light, and the polymerization rate and degree can be easily controlled by selecting the strength and wavelength of excitation. Photochemical polymerization can be proceeded even at relatively-low temperatures because its  
20 polymerization-initiating-energy is relatively low.

- [0003] At present, owing to the above characteristics of photopolymerization, there are rapidly arising demands for photopolymerizable compositions, which can be polymerized by irradiating visible light such as argon ion lasers, helium/neon lasers, and second harmonic of YAG lasers, with the development of new information recordings such as plates for printings and holographs. However, since conventional polymerizable compounds and polymerization initiators to be incorporated in photopolymerizable compositions selectively absorb ultraviolet rays, photopolymerizable  
25 compositions inevitably require photosensitizers as an essential technical factor when they are polymerized by irradiating visible light. Accordingly, in photopolymerizable compositions used in information recordings and electric equipments, polymerizable compounds should be used in combination with photosensitizers, polymerization initiators, binding resins, etc., which are all selected from various materials dependently on their uses. In general, methods comprising the steps of selecting materials for polymerizable compounds and/or polymerization initiators other than photosensitizers, and then screening photosensitizers which may appropriately sensitize the selected materials through trials  
30 and errors.

- [0004] The characteristics required in photosensitizers are a relatively-large molecular extinction coefficient in the visible region; an ability of sensitizing polymerizable compounds and polymerization initiators; a higher sensitization efficiency; a superior solubility in solvents and compatibility with other components; and a satisfactory stability. Such  
35 photosensitizers include, for example, merocyanine dyes as disclosed in Japanese Patent Kokai No. 151, 024/79, cyanine dyes as disclosed in Japanese Patent Kokai No. 29,803/83, stilbene derivatives as disclosed in Japanese Patent Kokai No. 56,403/84, coumarin derivatives as disclosed in Japanese Patent Kokai No. 23,901/88, pyran derivatives as disclosed in Japanese Patent Kokai No. 329,654/94, and methylene blue derivatives as disclosed in Japanese Patent  
40 Kokai No. 33,104/89. These photosensitizers, however, have unignorable disadvantages and remarkable advantages. There has been found no photosensitizer which consistently exerts the above characteristics in photo-polymerizable compositions containing appropriate materials.

[0005] Organic compounds sensitive to light, particularly, luminescent organic compounds are useful in the field of dye lasers and electroluminescence.

- 45 [0006] In the field of dye lasers, as disclosed, for example, by Kaoru IGARASHI in "Shikizai-Kyokai-shi", Vol. 70, No. 2, pp. 102-111 (1997), compounds which give luminescence in the visible region have been eagerly screened since the report of dye laser's lasing in 1960's. As the progress of information recording technology, there are arising demands for compounds which give luminescence in a longer wavelength region, more particularly, in the visible region.

- [0007] In the field of information displays, electro-luminescent devices (hereinafter abbreviated as "EL devices") are  
50 highlighted as displaying devices for the forthcoming generation. Nowadays, cathode-ray tubes are predominantly used in a relatively large size of information displaying means such as computer termini and TV receivers. The cathode-ray tubes are, however, relatively large in mass and weight and relatively high in operation voltage, thus the tubes are inadequate for consumer's equipments and small-sized equipments where portability is an important factor. In the small-sized equipments, display devices which have a depthless, light and plain form; and operate at a relatively-low operation voltage and wattage, are required. Because of the advantageous characteristics of a relatively-low operation voltage  
55 and wattage, liquid crystal devices are now widely used in various fields. However, because the contrast changes depending on view angles in the information displaying means using the liquid crystal devices, informations can be clearly displayed only when viewed within a specific view angle and the wattage is not effectively reduced as much as

expected because they generally need backlight. A device, which is now focused as a displaying device for overcoming the drawbacks as described above, is an organic electro-luminescent device or an organic EL device.

[0008] Organic EL device is a luminescent device which is formed by inserting a thin layer containing a luminescent agent between a cathode and an anode, and which utilizes luminescence such as fluorescence or phosphorescence emitted by the luminescent agent when a dc voltage is first supplied between the cathode and the anode to supply positive holes and electrons to the thin layer and to rebind the positive holes and the electrons in order to bring the luminescent agent into an excited state, and then the excited agent returns to the ground state. As a characteristic, organic EL device is appropriately changeable its luminous color tint by selecting appropriate host luminescent agents and changing guest luminescent agents as dopants used in combination with the host luminescent agents. Dependently on the combinations of the host and guest luminescent agents, the brightness and the life expectancy of luminescence can be improved by a large margin. It is said that organic EL device is a theoretically-excellent luminescent device, because it gives luminescence in itself and information displaying means therewith are free of view angles and low in energy consumption because it needs no backlight.

[0009] Although, in organic EL device which gives luminescence in green region, there has been reported an improvement of luminescent efficiency by incorporating guest luminescent agents, but it has been found no guest luminescent agent which effectively allows organic EL device to give luminescence in red region and it still remains far from emitting a pure red luminescence, poor in life expectancy, and insufficient in both durability and reliability. For example, the organic EL devices disclosed in Japanese Patent Kokai No. 6,042/98 and United States Patent No. 4,769,292 are insufficient in brightness and impure in red-color emission, and hence they have still problems in realizing full-color emission.

[0010] To supply organic EL devices at a lower cost, it is essential to find luminescent agents which do not intrinsically need doping by guest luminescent agents, make them simple the systematic structure of the organic EL devices and facilitating the step of vapor deposition in their preparations. Although various proposals have been made on luminescent agents in organic EL devices, there have been found no compound which fulfills the requirements as described above.

#### DISCLOSURE OF INVENTION

[0011] In view of the foregoing, the objects of the present invention are to provide organic compounds which absorb and/or emit lights in the visible region as well as to provide their uses in photochemical polymerization, organic EL devices, and dye lasers.

[0012] To solve the above objects, the present inventors eagerly screened compounds and researched and found that the pyran derivatives, which are obtainable through the step of reacting a compound having 4H-pyran skeleton with a compound having 3-formylcoumarin skeleton, have an absorption maximum in the visible region and substantially absorb visible light, and therefore they are very useful in photochemical polymerization. They also found that the pyran derivatives obtainable through such a step is very useful in dye lasers and organic EL devices because the pyran derivatives have the luminescent maxima in the visible region and emit visible light when excited.

[0013] More particularly, the present inventors confirmed that the pyran derivatives of the present invention emit at a high efficiency visible light, particularly, a light in red region when used in compounds for composing organic EL devices, in particular, when used in materials for composing luminescent layers of organic EL devices. The present invention was made on the basis of the creation of novel pyran derivatives and the findings of their industrially-useful features.

#### BRIEF DESCRIPTION OF DRAWING

[0014]

FIG. 1 is a brief figure of one organic EL device according to the present invention.

FIG. 2 is a brief figure of another organic EL device according to the present invention.

FIG. 3 is a brief figure of the other organic EL device according to the present invention.


FIG. 4 is a brief figure of another organic EL device according to the present invention.

[0015] Throughout FIGs 1 to 4, reference numerals 1 to 6 represent a substrate, anode, hole injection/transportation layer, luminous layer, electron injection/transportation layer, and cathode, respectively.


#### BEST MODE FOR CARRYING OUT THE INVENTION

[0016] The present invention relates to pyran derivatives represented by either of the following Formulae 1 and 2:

Formula 1



Formula 2



[0017] In Formulae 1 and 2,  $R_1$  to  $R_4$  independently represent a straight- or branched-chain of alkyl, alkenyl or aromatic hydrocarbon group which may contain one or more substituents. Dependently on uses, when  $R_1$  to  $R_4$  are alkyl groups, their chain lengths are usually selected from those up to 20 carbon atoms, desirably, 1 to 18 carbon atoms. Such alkyl groups include, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, n-pentyl, isopentyl, neopentyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, n-hexyl, isohexyl, cyclohexyl, 2-ethylhexyl, 5-ethylhexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, n-dodecyl, and n-octadecyl groups. Such alkenyl group include, for example, vinyl, 1-propenyl, 2-propenyl, isopropenyl, 2-butenyl, and 1, 3-butadienyl groups; Such aromatic hydrocarbon groups include, for example, phenyl, tolyl, xylyl, biphenyl, naphthyl, anthryl, and phenanthryl groups. Such substituents for these alkyl, alkenyl and aromatic hydrocarbon groups include, for example, halogens such as fluorine, chlorine, bromine, and iodine; alkoxy groups such as methoxy, ethoxy, propoxy, and butoxy groups; alkoxycarbonyl groups such as alkylcarboxy groups, methoxycarbonyl, ethoxycarbonyl groups, etc; carboxy, sulfonyl, alkylsulfonyl, aminosulfonyl, hydroxy, aromatic carboxy, cyano, phenylmethyl, 4-butylphenylmethyl, 4-butoxyphenylmethyl, 2-hydroxyethyl, 2-cyanoethyl, 2-carboxyethyl, 2-butoxyethyl, 2-(2-ethoxy)ethoxyethyl, 3-sulfoxypropyl, 4-sulfoxybutyl, 6-bromohexyl, and cycloalkyl groups.  $R_1$  and  $R_2$ , or  $R_3$  and  $R_4$  may form a five- or six-membered cyclic structure in cooperation with the nitrogen atom bonded thereunto or the benzene ring bonded to the nitrogen atom.


[0018] In Formulae 1 and 2,  $R_5$  and  $R_6$  independently represent cyano or alkyl halide; substituents such as ester, acyl and amido groups which are derived from carboxy group or carbonic acid; or monocyclic or heterocyclic groups which comprise one or more hetero atoms such as nitrogen, oxygen, sulfur, and selenium. Such heterocyclic groups include, for example, thiazolyl, benzothiazolyl, naphthothiazolyl, benzoxazolyl, naphthoxazolyl, imidazolyl, benzimidazolyl, naphthimidazolyl, and triazolyl groups. These heterocyclic groups may have one or more substituents such as halogens, alkyl, alkoxy, aromatic hydrocarbons, carboxy, sulfonic acid, organic acid esters, nitro, cyano, and alkyl halides such as trifluoroalkyl group.

[0019] Furthermore, in Formulae 1 and 2,  $R_7$  and  $R_8$  independently represent hydrogen; halogens such as fluoro, chloro, bromo, or iodo; cyano group; alkyl or halogenated alkyl group having up to six carbon atoms; or substituents derivable from carboxy group or carbonic acid, such as alkoxycarbonyl, acyl, and amido groups; where the alkyl group may have a straight-, cyclic-, or branched-chain structure or a substituent such as a halogen. In Formulae 1 and 2,  $R_9$


to R<sub>11</sub> independently represent hydrogen or lower alkyl group such as methyl, ethyl, or propyl group. The present invention includes *cis/trans* isomers of the pyran derivatives represented by Formula 1 or 2.

[0020] Concrete examples of the pyran derivatives represented by Formulae 1 and 2 are the compounds represented by the following Chemical Formulae 1 to 55. All of these pyran derivatives have absorption maxima and substantially absorb visible light, therefore they can be advantageously used in photosensitizers to photochemically polymerize polymerizable compounds. The pyran derivatives, which have luminescent maxima in visible region and emit visible light when excited, are very useful as laser-active substances in dye lasers, organic compounds which compose organic EL devices, particularly, host luminescent agents which form luminescent layers of organic EL devices, and guest luminescent agents to improve their luminescent efficiency and luminescent spectra by doping other host luminescent agents.


Chemical Formula 1:




Chemical Formula 2:




Chemical Formula 3:




Chemical Formula 4:




Chemical Formula 5:




Chemical Formula 6:




Chemical Formula 7:




Chemical Formula 8:




Chemical Formula 9:



Chemical Formula 10:




Chemical Formula 11:







Chemical Formula 12:




Chemical Formula 13:




Chemical Formula 14:




Chemical Formula 15:




Chemical Formula 16:




Chemical Formula 17:




Chemical Formula 18:




Chemical Formula 19:




Chemical Formula 20:




Chemical Formula 21:




Chemical Formula 22:




Chemical Formula 23:




Chemical Formula 24:




Chemical Formula 25:




Chemical Formula 26:




Chemical Formula 27:




Chemical Formula 28:




Chemical Formula 29:




Chemical Formula 30:




Chemical Formula 31:




Chemical Formula 32:




Chemical Formula 33:



Chemical Formula 34:




Chemical Formula 35:







Chemical Formula 36:




Chemical Formula 37:




Chemical Formula 38:




Chemical Formula 39:




Chemical Formula 40:




Chemical Formula 41:




Chemical Formula 42:




Chemical Formula 43:




Chemical Formula 44:




Chemical Formula 45:




Chemical Formula 46:




Chemical Formula 47:




Chemical Formula 48:




Chemical Formula 49:




Chemical Formula 50:



Chemical Formula 51:



Chemical Formula 52:




Chemical Formula 53:

5

10

15




Chemical Formula 54:

20

25

30




35

Chemical Formula 55:

40

45



50

[0021] The pyran derivatives of the present invention can be prepared by various methods. The following methods can be preferably used in economical view point: A method to produce the pyran derivative represented by Formula 1 through the step of reacting a compound represented by Formula 3 having  $R_1$ ,  $R_2$ ,  $R_5$ ,  $R_6$ ,  $R_{10}$  and  $R_{11}$  in Formula 1 with a compound represented by Formula 4 having  $R_3$ ,  $R_4$ ,  $R_7$  and  $R_9$  in Formula 1 in an appropriate solvent in the presence of a basic compound such as piperidine, pyridine, N,N-dimethylaniline, or triethylamine; and a method to produce the pyran derivative represented by Formula 2 through either a step of reacting a compound represented by Formula 5, having  $R_1$ ,  $R_2$ ,  $R_5$ ,  $R_6$ ,  $R_8$  and  $R_{10}$  in Formula 2, with a compound represented by Formula 4 having  $R_3$ ,  $R_4$ ,  $R_7$  and  $R_9$

in Formula 2 in an appropriate solvent in the presence of a basic compound such as piperidine, pyridine, N,N-dimethylaniline, or triethylamine; or a step of reacting a compound represented by Formula 6, having R<sub>5</sub> and R<sub>6</sub> in Formula 2, with a compound represented by Formula 4 having R<sub>3</sub>, R<sub>4</sub>, R<sub>7</sub> and R<sub>9</sub> in Formula 2 and a compound represented by Formula 7 having R<sub>1</sub>, R<sub>2</sub>, R<sub>8</sub> and R<sub>10</sub> in Formula 2 in an appropriate solvent in the presence of a similar basic compound as used above. In the method to react compounds represented by Formula 6 with compounds represented by Formulae 4 and 7, the pyran derivatives of the present invention, which bind the same substituents at the C-2 and C-6 positions of 4H-pyran ring, are obtained when single compounds are used as compounds represented by Formulae 4 and 7. The pyran derivatives represented by chemical Formulae 1 to 55 can be prepared by these methods.

10

15

20

25

30

35


40

45


50

55


Formula 3:




Formula 4:



Formula 5:




Formula 6:





Formula 7:



[0022] The pyran derivatives thus obtained can be used intact in a reaction mixture form dependently on uses, however, prior to use, they are usually purified by conventional methods generally used in the purification of analogous compounds, for example, separation, decantation, filtration, concentration, thin-layer chromatography, column chromatography, gas chromatography, high-performance liquid chromatography, distillation, sublimation, and crystallization, which can be used in combination, if necessary. When used in photosensitizers, the pyran derivatives would preferably be at least distilled and/or crystallized prior to use, while when used in laser-active substances for laser oscillating devices and in luminescent agents for organic EL devices, the pyran derivatives would preferably be further purified, for example, by a method such as sublimation.

[0023] As described above, since the pyran derivatives of the present invention substantially absorb visible light, particularly, longer wavelength of light, and have characteristics that they sensitize various polymerizable compounds and polymerization initiators, they can be very useful as photosensitizers to polymerize photochemically the polymerizable compounds by using light with longer wavelengths than ultraviolet ray, particularly, by using visible light. In use, the pyran derivatives of the present invention are usually preparable into compositions in combination with polymerizable compounds, polymerization initiators, and/or binding resins. However, depending on the kinds of the pyran derivatives and the final use of photopolymerizable compounds, polymerization initiators, and/or binding resins can be omitted.

[0024] The polymerizable compounds, to which the photosensitizers of the present invention can be applied, are, for example, monomers, oligomers, prepolymers, and mixtures thereof, which contain at least one polymerizable multiple bond such as an ethylenic-double-bond. Such polymerizable compounds include, for example, ethylacrylate, hydroxyethylacrylate, ethylene glycol dimethacrylate, pentaerythritol triacrylate, pentaerythritol trimethacrylate, polyester methacrylate, polyurethane methacrylate, and epoxymethacrylate. However, the polymerizable compounds, which are incorporated into the photopolymerizable compositions of the present invention, should not be restricted to the above compounds and include all polymerizable compounds which can be photochemically polymerized by using the photosensitizers of the present invention.

[0025] Examples of the polymerization initiators are organic peroxides such as di-t-butylperoxyisophthalate, 3,3',4,4'-tetrakis(t-butylperoxy) benzophenone, ethylmethylketone, 2,5-dimethyl-2,5-bis (t-butylperoxy)-3-hexane, di-t-butylperoxide, 2, 5-bis (hydroperoxy)-2,5-dimethylhexane, t-butylhydroperoxide, butyl-4,4-bis (t-butylperoxy) valylate, and 1,1-bis (t-butylperoxy)-3,3,5-trimethylcyclohexane; halogenated hydrocarbons such as 2,4,6-trichloromethyl-s-triazine; bisimidazol; benzoylalkylether; iron-allene complexes; titanocene compounds; N-phenylglycine, and diphenyliodonium salts, which are polymerization initiators conventionally used in photopolymerization and which can be used in combination, if necessary.

[0026] Dependently on uses, any binding resins, which are generally used in photopolymerizable compositions, can be used in the present invention. Such binding resins are, for example, poly-N-vinylpyrrolidone, poly (vinyl acetate), poly (vinyl butyral), poly (vinyl carbazole), polystyrene, poly (methyl methacrylate), poly (ethylene oxide), poly (butyl methacrylate), styrene-maleic acid ester, poly (methyl methacrylate)-methacrylic acid, and poly-N-vinylpyrrolidone-glycidyl methacrylate.

[0027] The photosensitizers of the present invention include one or more of the pyran derivatives of the present invention. Usually, to obtain photopolymerizable compositions by using photosensitizers of the present invention, they can be prepared by incorporating into one part by weight of any of the photosensitizers of the present invention 1-1,000 parts by weight, and preferably 10-500 parts by weight of any of the aforesaid polymerizable compounds; and if necessary, further incorporating into the mixtures any of the aforesaid binding resins in an amount up to 1,000 parts by weight, and preferably up to 500 parts by weight, and further incorporating into the resulting mixtures any of the aforesaid polymerization initiators in an amount of 0.1-10 parts by weight, and preferably 0.5-5 parts by weight. In addition, the following quinonoid or phenolic thermopolymerization inhibitors such as hydroquinone, pyrogallol, and 2,6-di-t-

butyl-p-cresol; and plasticizers such as saturated- or unsaturated-carboxylic esters including phthalic acid esters and adipic acid esters, colorants, preservatives, stabilizers, surface protectants, smoothers, and coating aids can be appropriately added to the above photopolymerizable compositions.

**[0028]** Usually, the photopolymerizable compositions of the present invention can be used by dissolving in appropriate solvents into solutions, coating the solutions over supporting means appropriately, and drying the coated means. As the solvents, the followings can be used: Hydrocarbons such as pentane, hexane, cyclohexane, petroleum ether, petroleum benzine, isooctane, benzene, toluene, and xylene; halides such as carbon tetrachloride, chloroform, 1,2-dichloroethane, 1,2-dibromoethane, trichloroethylene, tetrachloroethylene, chlorobenzene, and  $\alpha$ -dichlorobenzene; alcohols and phenols such as methanol, ethanol, 1-propanol, 2-propanol, 2,2,2-trifluoroethanol, 1-butanol, 2-butanol, isobutyl alcohol, isopentyl alcohol, cyclohexanol, ethylene glycol, propylene glycol, 2-methoxyethanol, 2-ethoxyethanol, phenol, benzyl alcohol, cresol, diethylene glycol, triethylene glycol, and glycerin; ethers such as diethyl ether, diisopropyl ether, tetrahydrofuran, tetrahydropyran, 1,4-dioxane, anisole, 1,2-dimethoxyethane, diethylene glycol dimethyl ether, dicyclohexyl-18-crown-6, methylcarbitol, ethylcarbitol, propylene glycol monomethyl ether, and propylene glycol monomethyl ether acetate; ketones such as acetone, ethyl methyl ketone, and cyclohexanone; esters such as ethyl acetate, butyl acetate, amyl acetate, ethyl propionate, ethylene carbonate, propylene carbonate, and phosphoric trimethyl; amides such as formamide, N-methylformamide, N,N-dimethylformamide, N-methylacetamide, N,N-dimethylacetamide, and hexamethyl phosphoric triamide; nitriles such as acetonitrile, propionitrile, succinonitrile, and benzonitrile; nitro compounds such as nitromethane and nitrobenzene; amines such as ethylenediamine, pyridine, piperidine, and morpholine; and sulfur-containing compounds such as dimethylsulfoxide and sulfolane. These solvents can be arbitrarily used in combination, if necessary.

**[0029]** Any support means generally used in this field can be used in the present invention. Such support means are, for example, metals such as aluminum, magnesium, copper, zinc, chromium, nickel, and iron; alloy sheets thereof; papers such as high-quality papers, art papers, and exfoliative papers; inorganic sheets such as glasses and ceramics; and plastic sheets such as polyethylene phthalate, polyethylene, poly(methyl methacrylate), poly(vinyl chloride), copolymer of vinyl chloride-vinylidene chloride, polystyrene, nylon, cellulose acetate, and cellulose acetate butyrate.

**[0030]** The polymerization method used in the present invention is not specifically restricted and, for example, it can be photoinitiation polymerization where light participates only in the initiation stage of radical-, ion-, and ring opening-polymerizations for polymerizable compounds; or may be photo-polyaddition polymerization where light participates in the propagation stage of polymerizable compounds. The exposure illuminants usable in the present invention are conventional illuminants, for example, visible light with wavelengths of more than 400 nm such as sun light, carbon arcs, mercury lamps, xenon lamps, metalhalide lamps, luminescent lamps, tungsten lamps, argon ion lasers, krypton ion lasers, helium-cadmium lasers, helium-neon lasers, and the second harmonic wavelength by YAG laser.

**[0031]** The present invention also provides the uses as luminous agents of the pyran derivatives in organic EL devices. Since most of the pyran derivatives of the present invention have luminescent maxima such as fluorescent maxima in the visible region at wavelengths of more than 600 nm, particularly, in the red region at wavelengths of 610 to 730 nm, the pyran derivatives can be advantageously useful as luminous agents for organic EL devices.

**[0032]** The luminous agents for organic EL devices of the present invention can be advantageously used both in monolayer- and laminate-types of organic EL devices. The action of the organic EL devices essentially comprises the step of injecting electrons and holes from an electrode, allowing the electrons and the holes to transport through solid members, forming singlet or triplet excitons by recombining the electrons and the holes, and allowing the excitons to emit. These steps do not intrinsically differ both in the monolayer- and the laminate-types of organic EL devices. However, in general, the laminate-type organic EL devices attain desirable properties easier than those of the monolayer-type because the functions required in each step can be allowed to share by plural materials and optimized separately for each material in the laminate-type organic EL devices, while in the monolayer-type organic EL devices the properties of the aforesaid four steps can be improved only by changing the molecular structure of luminescent agents.


**[0033]** Now explaining the organic EL devices of the present invention with reference to a laminate-type organic EL device, the organic EL device, as shown in FIG. 1, is essentially composed of anode 2 to be energized with positive voltage, cathode 6 to be energized with negative voltage, and luminous layer 4 to give luminescence by recombining holes and electrons. The organic EL device may have hole injection/transportation layer 3 which accelerates the injection and transportation of holes from anode 2, and/or electron injection/transportation layer 5 which accelerates the injection and transportation of electrons from cathode 6, as shown in FIGs. 2 to 4, if necessary. In FIGs. 1 to 4, the reference numeral 1 represents a substrate, i.e., a substrate in general including glasses such as soda glass, barium boron silicate glass, and aluminosilicate glass; plastics; or ceramics. A preferable substrate is a transparent glass or plastic, and an opaque ceramic such as silicon can be used in combination with a transparent electrode.

**[0034]** The reference numeral 2 represents an anode which is usually formed by attaching closely to either surface of substrate 1 by a method such as vacuum deposition, molecular beam deposition, chemical deposition (CDV), immersion, spin-coating, bar-coating, roll-coating, casting, or spattering, and then coating thereupon a metal or an electrically conductive material, which is electrically low in resistivity and high in transmissivity over visible region, into a thin layer


having a thickness of 10-1,000 nm, desirably, 50-500 nm to give a resistivity of anode 2  $1\text{k}\Omega/\square$ . The following compounds can be used in the electric conductive materials in the present invention: Metals such as gold, platinum, aluminum, and nickel; metal oxides such as tin oxide, indium oxide, and mixture systems of tin oxide and indium oxide (hereinafter abbreviated as "ITO"); and electric conductive oligomers or polymers which have aniline, thiophene or pyrrole as a repeating unit. Among these, ITO has a characteristic of being easily processed into low-resistive ones and etched with acids to easily form minute patterns.

[0035] The reference numeral 3 represents a hole injection/transportation layer which is usually formed by attaching closely to anode 2 by the same methods as used in anode 2 to make a material for hole injection/transportation layer into a thin layer having a thickness of 1-1,000 nm. In the hole injection/transportation layer, desirably used are materials which give a hole mobility of at least  $10^{-6}\text{ cm}^2/\text{V} \cdot \text{sec}$  under electric fields of  $10^4$  to  $10^6\text{ V/cm}$  and a relatively-low ionized electric potential. Such materials include, for example, aromatic tertiary amine, styryl amine, triazole derivatives, pyrazoline derivatives, pyrazolone derivatives, phenylenediamine derivatives, allylamine derivatives, amine substituted chalcone derivatives, oxazole derivatives, styryl anthracene derivatives, fluorenone derivatives, hydrazone derivatives, stilbene derivatives, and so-called "star burst type molecules", which all can be arbitrarily used in combination, if necessary. The materials for hole injection/transportation layer include, for example, bis [di (p-tolyl) aminophenyl] -1,1-cyclohexane, N,N'-diphenyl-N,N'-bis(3-methylphenyl)-1,1'-biphenyl-4,4'-diamine, and N,N'-diphenyl-N,N'-bis (1-naphthyl)-1,1'-biphenyl-4,4'-diamine.


Chemical Formula 56:




Chemical Formula 57:




Chemical Formula 58:




Chemical Formula 59:



Chemical Formula 60:




Chemical Formula 61:




[0036] The reference numeral 4 represents luminous layer which is usually formed by attaching closely to hole injection/transportation layer 3, and then, by the same method as used in anode 2, coating thereupon the pyran derivative of the present invention alone or in combination with an appropriate host luminescent agent to form a thin monolayer or separated two-layers having, usually, a thickness of 1-1,000 nm, preferably, 10-200 nm. Though the pyran derivatives of the present invention *per se* exert high luminous efficiency, they effectively suppress the quenching of concentration of host luminescent agents and exert a higher luminous efficiency when doped in an appropriate amount to the host luminescent agents. The doping amount of the pyran derivatives of the present invention is usually 0.1-30 % by weight, desirably, 1-10 % by weight, and it may be varied depending on the kind of the host luminescent agents. The host luminescent agents used in combination with the pyran derivatives of the present invention are, for example, the following compounds represented by Formulae 8 to 13.


Formula 8:




Formula 9:




Formula 10:




Formula 11:



Formula 12:



Formula 13:



**[0037]** In Formula 8, all of  $R_{12}$  to  $R_{14}$  independently represent an aromatic hydrocarbon group or aromatic heterocyclic group, which may contain a substituent and bind each other to form a cyclic structure.

**[0038]** In Formula 9, all of  $R_{15}$  to  $R_{18}$  independently represent an aromatic hydrocarbon group or aromatic heterocyclic group, which may contain a substituent and bind each other to form a cyclic structure. Y represents an arylene group which may contain a substituent.

**[0039]** In Formula 10, all of  $R_{19}$  to  $R_{24}$  independently represent hydrogen or a substituent selected from hydroxy, nitro, cyano, carboxy, amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxy carbonyl groups, where the amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxy carbonyl group may contain a substituent.  $L_1$  represents alkyl, alkenyl, cycloalkyl, aromatic hydrocarbon, aromatic heterocyclic, or aralkyl group, where the alkyl, alkenyl, cycloalkyl, aromatic hydrocarbon, aromatic heterocyclic, or aralkyl group may contain a substituent. m represents an integer selected from 0 to 2, and n represents an integer selected from 1 to 3. M represents a metal ion with a valency of (n+m).

**[0040]** In Formula 11, all of  $R_{25}$  to  $R_{36}$  independently represent hydrogen or a substituent selected from halogens, hydroxy, nitro, cyano, carboxy, amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxy carbonyl groups, where the amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxy carbonyl groups may contain a substituent and bind each other to form a cyclic structure.  $L_2$  represents alkylene, alkenylene, cycloalkylene, bivalent aromatic hydrocarbon, or bivalent aromatic heterocyclic group, where the alkylene, alkenylene, cycloalkylene, aromatic hydrocarbon, or aromatic heterocyclic group may contain a substituent. 1 represents an integer of 0 or 1 and s represents an integer of 1 or 2. M represents a metal ion with a valency of (s+1).

**[0041]** In Formula 12, all of  $R_{37}$  to  $R_{40}$  independently represent cyano, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, or aryloxy group, where the alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, or aryloxy group may contain a substituent and bind each other to form a cyclic structure. Y represents an arylene group, which may contain a substituent.

**[0042]** In Formula 13, all of  $R_{41}$  to  $R_{46}$  independently represent an aromatic hydrocarbon group or aromatic heterocyclic group, which may contain one or more substituents or the substituents may bind each other to form a cyclic structure. Z represents a trivalent aromatic hydrocarbon group or trivalent aromatic heterocyclic group, which may contain one or more substituents.

**[0043]** Explaining the substituents in Formulae 8 to 13, the halogens can be fluoro, chloro, bromo, or iodo. A substituted or unsubstituted amino group is represented by  $-NX_1X_2$ , and  $X_1$  and  $X_2$  are each independently hydrogen, methyl, ethyl, propyl, isopropyl, n-butyl, s-butyl, isobutyl, t-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxyisobutyl, 1,2-dihydroxyethyl, 1,3-dihydroxyisopropyl, 2,3-dihydroxy-t-butyl, 1,2,3-trihydroxypropyl, chloromethyl, 1-chloroethyl, 2-chloroethyl, 2-chloroisobutyl, 1,2-dichloroethyl, 1,3-dichloroisopropyl, 2,3-dichloro-t-butyl, 1,2,3-trichloropropyl, bromomethyl, 1-bromoethyl, 2-bromoethyl, 2-bromoisobutyl, 1,2-dibromoethyl, 1,3-dibromoisopropyl, 2,3-dibromo-t-butyl, 1,2,3-tribromopropyl, iodomethyl, 1-iodoethyl, 2-iodoethyl, 2-iodoisobutyl, 1,2-diiodoethyl, 1,3-diiodoisopropyl, 2,3-diiodo-t-butyl, 1,2,3-triiodopropyl, aminomethyl, 1-aminoethyl, 2-aminoethyl, 2-aminoisobutyl, 1,2-diaminoethyl, 1,3-diaminoisopropyl, 2,3-diamino-t-butyl, 1,2,3-triaminopropyl, cyanomethyl, 1-cyanoethyl, 2-cyanoethyl, 2-cyanoisobutyl, 1,2-dicyanoethyl, 1,3-dicyanoisopropyl, 2,3-dicyano-t-butyl, 1,2,3-tricyanopropyl, nitromethyl, 1-nitroethyl, 2-nitroethyl, 2-nitroisobutyl, 1,2-dinitroethyl, 1,3-dinitroisopropyl, 2,3-dinitro-t-butyl, 1,2,3-trinitropropyl, phenyl, 1-naphthyl, 2-naphthyl, 1-anthryl, 2-anthryl, 9-anthryl, 1-phenanthryl, 2-phenanthryl, 3-phenanthryl, 4-phenanthryl, 9-phenanthryl, 1-naphthacenyl, 2-naphthacenyl, 9-naphthacenyl, 4-styrylphenyl, 1-pyrenyl, 2-pyrenyl, 4-pyrenyl, 2-biphenyl, 3-biphenyl, 4-biphenyl, p-terphenyl-4-yl, p-terphenyl-3-yl, p-terphenyl-2-



yl, m-terphenyl-4-yl, m-terphenyl-3-yl, m-terphenyl-2-yl, o-tolyl, m-tolyl, p-tolyl, p-t-butylphenyl, p-(2-phenylpropyl)phenyl, 3-methyl-2-naphthyl, 4-methyl-1-naphthyl, 4-methyl-1-anthryl, 4'-methylbiphenyl, 4"-t-butyl-p-terphenyl-4-yl, 2-pyrrolyl, 3-pyridyl, pyrazinyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl, 1-isoidolyl, 3-isoidolyl, 4-isoidolyl, 5-isoidolyl, 6-isoidolyl, 7-isoidolyl, 2-furyl, 3-furyl, 2-benzofuryl, 3-benzofuryl, 4-benzofuryl, 5-benzofuryl, 6-benzofuryl, 7-benzofuryl, 1-isobenzofuryl, 3-isobenzofuryl, 4-isobenzofuryl, 5-isobenzofuryl, 6-isobenzofuryl, 7-isobenzofuryl, 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl, 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 6-quinoxalyl, 1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl, 1-phenanthridinyl, 2-phenanthridinyl, 3-phenanthridinyl, 4-phenanthridinyl, 6-phenanthridinyl, 7-phenanthridinyl, 8-phenanthridinyl, 9-phenanthridinyl, 10-phenanthridinyl, 1-acridinyl, 2-acridinyl, 3-acridinyl, 4-acridinyl, 9-acridinyl, 1,7-phenanthroline-2-yl, 1,7-phenanthroline-3-yl, 1,7-phenanthroline-4-yl, 1,7-phenanthroline-5-yl, 1,7-phenanthroline-6-yl, 1,7-phenanthroline-8-yl, 1,7-phenanthroline-9-yl, 1,7-phenanthroline-10-yl, 1,8-phenanthroline-2-yl, 1,8-phenanthroline-3-yl, 1,8-phenanthroline-4-yl, 1,8-phenanthroline-5-yl, 1,8-phenanthroline-6-yl, 1,8-phenanthroline-7-yl, 1,8-phenanthroline-9-yl, 1,8-phenanthroline-10-yl, 1,9-phenanthroline-2-yl, 1,9-phenanthroline-3-yl, 1,9-phenanthroline-4-yl, 1,9-phenanthroline-5-yl, 1,9-phenanthroline-6-yl, 1,9-phenanthroline-7-yl, 1,9-phenanthroline-8-yl, 1,9-phenanthroline-9-yl, 1,9-phenanthroline-10-yl, 1,10-phenanthroline-2-yl, 1,10-phenanthroline-3-yl, 1,10-phenanthroline-4-yl, 1,10-phenanthroline-5-yl, 2,9-phenanthroline-1-yl, 2,9-phenanthroline-3-yl, 2,9-phenanthroline-4-yl, 2,9-phenanthroline-5-yl, 2,9-phenanthroline-6-yl, 2,9-phenanthroline-7-yl, 2,9-phenanthroline-8-yl, 2,9-phenanthroline-10-yl, 2,8-phenanthroline-1-yl, 2,8-phenanthroline-3-yl, 2,8-phenanthroline-4-yl, 2,8-phenanthroline-5-yl, 2,8-phenanthroline-6-yl, 2,8-phenanthroline-7-yl, 2,8-phenanthroline-9-yl, 2,8-phenanthroline-10-yl, 2,7-phenanthroline-1-yl, 2,7-phenanthroline-3-yl, 2,7-phenanthroline-4-yl, 2,7-phenanthroline-5-yl, 2,7-phenanthroline-6-yl, 2,7-phenanthroline-8-yl, 2,7-phenanthroline-9-yl, 2,7-phenanthroline-10-yl, 1-phenazinyl, 2-phenazinyl, 1-phenothiazinyl, 2-phenothiazinyl, 3-phenothiazinyl, 4-phenothiazinyl, 1-phenoxazinyl, 2-phenoxazinyl, 3-phenoxazinyl, 4-phenoxazinyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-oxadiazolyl, 5-oxadiazolyl, 3-furazanyl, 2-thienyl, 3-thienyl, 2-methylpyrrole-1-yl, 2-methylpyrrole-3-yl, 2-methylpyrrole-4-yl, 2-methylpyrrole-5-yl, 3-methylpyrrole-1-yl, 3-methylpyrrole-2-yl, 3-methylpyrrole-4-yl, 3-methylpyrrole-5-yl, 2-t-butylpyrrole-4-yl, 3-(2-phenylpropyl)pyrrole-1-yl, 2-methyl-1-indolyl, 4-methyl-1-indolyl, 2-methyl-3-indolyl, 4-methyl-3-indolyl, 2-t-butyl-1-indolyl, 4-t-butyl-1-indolyl, 2-t-butyl-3-indolyl, or 4-t-butyl-3-indolyl group.

**[0044]** The substituted or unsubstituted alkyl group is a straight- or branched-chain alkyl group, which may contain one or more substituents such as methyl, ethyl, propyl, isopropyl, n-butyl, s-butyl, isobutyl, t-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxyisobutyl, 1,2-dihydroxyethyl, 1,3-dihydroxyisopropyl, 2,3-dihydroxy-t-butyl, 1,2,3-trihydroxypropyl, chloromethyl, 1-chloroethyl, 2-chloroethyl, 2-chloroisobutyl, 1,2-dichloroethyl, 1,3-dichloroisopropyl, 2,3-dichloro-t-butyl, 1,2,3-trichloropropyl, bromomethyl, 1-bromoethyl, 2-bromoethyl, 2-bromoisobutyl, 1,2-dibromoethyl, 1,3-dibromoisopropyl, 2,3-dibromo-t-butyl, 1,2,3-tribromopropyl, iodomethyl, 1-iodoethyl, 2-iodoethyl, 2-iodoisobutyl, 1,2-diiodoethyl, 1,3-diiodoisopropyl, 2,3-diiodo-t-butyl, 1,2,3-triiodopropyl, aminomethyl, 1-aminoethyl, 2-aminoethyl, 2-aminoisobutyl, 1,3-diaminoethyl, 1,3-diaminoisopropyl, 2,3-diamino-t-butyl, 1,2,3-triaminopropyl, cyanomethyl, 1-cyanoethyl, 2-cyanoethyl, 2-cyanoisobutyl, 1,2-dicyanoethyl, 1,3-dicyanoisopropyl, 2,3-dicyano-t-butyl, 1,2,3-tricyanopropyl, nitromethyl, 1-nitroethyl, 2-nitromethyl, 2-nitroisobutyl, 1,2-dinitroethyl, 1,3-dinitroisopropyl, 2,3-dinitro-t-butyl, and 1,2,3-trinitropropyl groups.

**[0045]** The substituted or unsubstituted alkenyl group is a straight- or branched-chain alkenyl group, which may contain one or more substituents such as vinyl, allyl, 1-butenyl, 2-butenyl, 3-butenyl, 1,3-butadienyl, 1-methylvinyl, styryl, 2,2-diphenylvinyl, 1,2-diphenylvinyl, 1-methylallyl, 1,1-dimethylallyl, 2-methylallyl, 1-phenylallyl, 2-phenylallyl, 3-phenylallyl, 3,3-diphenylallyl, 1,2-dimethylallyl, 1-phenyl-1-butenyl, and 3-phenyl-1-butenyl groups.

**[0046]** The substituted or unsubstituted cycloalkyl group is a branched-chain cycloalkyl group, which may contain one or more substituents such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and 4-methylcyclohexyl groups.

**[0047]** The substituted or unsubstituted alkoxy group is represented by -OY, where Y is, for example, methyl, ethyl, propyl, isopropyl, n-butyl, s-butyl, isobutyl, t-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxyisobutyl, 1,2-dihydroxyethyl, 1,3-dihydroxyisopropyl, 2,3-dihydroxy-t-butyl, 1,2,3-trihydroxypropyl, chloromethyl, 1-chloroethyl, 2-chloroethyl, 2-chloroisobutyl, 1,2-dichloroethyl, 1,3-dichloroisopropyl, 2,3-dichloro-t-butyl, 1,2,3-trichloropropyl, bromomethyl, 1-bromoethyl, 2-bromoethyl, 2-bromoisobutyl, 1,2-dibromoethyl, 1,3-dibromoisopropyl, 2,3-dibromo-t-butyl, 1,2,3-tribromopropyl, iodomethyl, 1-iodoethyl, 2-iodoethyl, 2-iodoisobutyl, 1,2-diiodoethyl, 1,3-diiodoisopropyl, 2,3-diiodo-t-butyl, 1,2,3-triiodopropyl, aminomethyl, 1-aminoethyl, 2-aminoethyl, 2-aminoisobutyl, 1,2-diaminoethyl, 1,3-diaminoisopropyl, 2,3-diamino-t-butyl, 1,2,3-triaminopropyl, cyanomethyl, 1-cyanoethyl, 2-cyanoethyl, 2-cyanoisobutyl, 1,2-dicyanoethyl, 1,3-dicyanoisopropyl, 2,3-dicyano-t-butyl, 1,2,3-tricyanopropyl, nitromethyl, 1-nitroethyl, 2-nitroethyl, 2-nitroisobutyl, 1,2-dinitroethyl, 1,3-dinitroisopropyl, 2,3-dinitro-t-butyl, or 1,2,3-trinitropropyl group.

**[0048]** The substituted or unsubstituted aromatic hydrocarbon group is a monovalent or multivalent aromatic hydrocarbon group, which may contain one or more substituents such as phenyl, 1-naphthyl, 2-naphthyl, 1-anthryl, 2-anthryl, 9-anthryl, 1-phenanthryl, 2-phenanthryl, 3-phenanthryl, 4-phenanthryl, 9-phenanthryl, 1-naphthacenyl, 2-naphthace-

nyl, 9-naphthacenyl, 1-pyrenyl, 2-pyrenyl, 4-pyrenyl, 2-biphenyl, 3-biphenyl, 4-biphenyl, p-terphenyl-4-yl, p-terphenyl-3-yl, p-terphenyl-2-yl, m-terphenyl-4-yl, m-terphenyl-3-yl, m-terphenyl-2-yl, o-tolyl, m-tolyl, p-tolyl, p-t-butylphenyl, p-(2-phenylpropyl) phenyl, 3-methyl-2-naphthyl, 4-methyl-1-naphthyl, 4-methyl-1-anthryl, 4'-methylbiphenyl, and 4"-t-butyl-p-terphenyl-4-yl groups.

5 **[0049]** The substituted or unsubstituted aromatic heterocyclic group is a monovalent or multivalent group of monocyclic or polycyclic heterocyclic compound which may contain one or more heteroatoms such as nitrogen, oxygen, sulfur, and selenium, where the aromatic heterocyclic group may contain one or more substituents. Examples of such an aromatic heterocyclic group are 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, pyrazinyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl, 1-isindolyl, 2-isindolyl, 3-isindolyl, 4-isindolyl, 5-isindolyl, 6-isindolyl, 7-isindolyl, 2-furyl, 3-furyl, 2-benzofuryl, 3-benzofuryl, 4-benzofuryl, 5-benzofuryl, 6-benzofuryl, 7-benzofuryl, 1-isobenzofuryl, 3-isobenzofuryl, 4-isobenzofuryl, 5-isobenzofuryl, 6-isobenzofuryl, 7-isobenzofuryl, 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl, 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 6-quinoxalyl, 1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl, 9-carbazolyl, 1-phenanthridinyl, 2-phenanthridinyl, 3-phenanthridinyl, 4-phenanthridinyl, 6-phenanthridinyl, 7-phenanthridinyl, 8-phenanthridinyl, 9-phenanthridinyl, 10-phenanthridinyl, 1-acridinyl, 2-acridinyl, 3-acridinyl, 4-acridinyl, 9-acridinyl, 1,7-phenanthroline-3-yl, 1,7-phenanthroline-4-yl, 1,7-phenanthroline-5-yl, 1,7-phenanthroline-6-yl, 1,7-phenanthroline-8-yl, 1,7-phenanthroline-9-yl, 1,7-phenanthroline-10-yl, 1,8-phenanthroline-2-yl, 1,8-phenanthroline-3-yl, 1,8-phenanthroline-4-yl, 1,8-phenanthroline-5-yl, 1,8-phenanthroline-6-yl, 1,8-phenanthroline-7-yl, 1,8-phenanthroline-9-yl, 1,8-phenanthroline-10-yl, 1,9-phenanthroline-2-yl, 1,9-phenanthroline-3-yl, 1,9-phenanthroline-4-yl, 1,9-phenanthroline-5-yl, 1,9-phenanthroline-6-yl, 1,9-phenanthroline-7-yl, 1,9-phenanthroline-8-yl, 1,9-phenanthroline-10-yl, 1,10-phenanthroline-2-yl, 1,10-phenanthroline-3-yl, 1,10-phenanthroline-4-yl, 1,10-phenanthroline-5-yl, 2,9-phenanthroline-1-yl, 2,9-phenanthroline-3-yl, 2,9-phenanthroline-4-yl, 2,9-phenanthroline-5-yl, 2,9-phenanthroline-6-yl, 2,9-phenanthroline-7-yl, 2,9-phenanthroline-8-yl, 2,9-phenanthroline-10-yl, 2,8-phenanthroline-1-yl, 2,8-phenanthroline-3-yl, 2,8-phenanthroline-4-yl, 2,8-phenanthroline-5-yl, 2,8-phenanthroline-6-yl, 2,8-phenanthroline-7-yl, 2,8-phenanthroline-9-yl, 2,8-phenanthroline-10-yl, 2,7-phenanthroline-1-yl, 2,7-phenanthroline-3-yl, 2,7-phenanthroline-4-yl, 2,7-phenanthroline-5-yl, 2,7-phenanthroline-6-yl, 2,7-phenanthroline-8-yl, 2,7-phenanthroline-9-yl, 2,7-phenanthroline-10-yl, 1-phenazinyl, 2-phenazinyl, 1-phenothiazinyl, 2-phenothiazinyl, 3-phenothiazinyl, 4-phenothiazinyl, 10-phenothiazinyl, 1-phenoxazinyl, 2-phenoxazinyl, 3-phenoxazinyl, 4-phenoxazinyl, 10-phenoxazinyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-oxadiazolyl, 5-oxadiazolyl, 3-furazanyl, 2-thienyl, 3-thienyl, 2-methylpyrrole-1-yl, 2-methylpyrrole-3-yl, 2-methylpyrrole-4-yl, 2-methylpyrrole-5-yl, 3-methylpyrrole-1-yl, 3-methylpyrrole-2-yl, 3-methylpyrrole-4-yl, 3-methylpyrrole-5-yl, 2-t-butylpyrrole-4-yl, 3-(2-phenylpropyl) pyrrole-1-yl, 2-methyl-1-indolyl, 4-methyl-1-indolyl, 2-methyl-3-indolyl, 4-methyl-3-indolyl, 2-t-butyl-1-indolyl, 4-t-butyl-1-indolyl, 2-t-butyl-3-indolyl, or 4-t-butyl-3-indolyl group.

35 **[0050]** Examples of the substituted or unsubstituted aralkyl group are benzyl, 1-phenylethyl, 2-phenylethyl, 1-phenylisopropyl, 2-phenylisopropyl, phenyl-t-butyl,  $\alpha$ -naphthylmethyl, 1- $\alpha$ -naphthylethyl, 2- $\alpha$ -naphthylethyl, 1- $\alpha$ -naphthylisopropyl, 2- $\alpha$ -naphthylisopropyl,  $\beta$ -naphthylmethyl, 1- $\beta$ -naphthylethyl, 2- $\beta$ -naphthylethyl, 1- $\beta$ -naphthylisopropyl, 2- $\beta$ -naphthylisopropyl, 1-pyrrolylmethyl, 2-(1-pyrrolyl) ethyl, p-methylbenzyl, m-methylbenzyl, o-methylbenzyl, p-chlorobenzyl, m-chlorobenzyl, o-chlorobenzyl, p-bromobenzyl, m-bromobenzyl, o-bromobenzyl, p-iodobenzyl, m-iodobenzyl, o-iodobenzyl, p-hydroxybenzyl, m-hydroxybenzyl, o-hydroxybenzyl, p-aminobenzyl, m-aminobenzyl, o-aminobenzyl, p-nitrobenzyl, m-nitrobenzyl, o-nitrobenzyl, p-cyanobenzyl, m-cyanobenzyl, o-cyanobenzyl, 1-hydroxy-2-phenylisopropyl, and 1-chloro-2-phenylisopropyl groups.

45 **[0051]** The substituted or unsubstituted aryloxy group is represented by -OZ, where Z is, for example, phenyl, 1-naphthyl, 2-naphthyl, 1-anthryl, 2-anthryl, 9-anthryl, 1-phenanthryl, 2-phenanthryl, 3-phenanthryl, 4-phenanthryl, 9-phenanthryl, 1-naphthacenyl, 2-naphthacenyl, 9-naphthacenyl, 1-pyrenyl, 2-pyrenyl, 4-pyrenyl, 2-biphenyl, 3-biphenyl, 4-biphenyl, p-terphenyl-4-yl, p-terphenyl-3-yl, p-terphenyl-2-yl, m-terphenyl-4-yl, m-terphenyl-3-yl, m-terphenyl-2-yl, o-tolyl, m-tolyl, p-tolyl, p-t-butylphenyl, p-(2-phenylpropyl) phenyl, 3-methyl-2-naphthyl, 4-methyl-1-naphthyl, 4-methyl-1-anthryl, 4'-methylbiphenyl, 4"-t-butyl-p-terphenyl-4-yl, 2-pyrrolyl, 3-pyrrolyl, pyrazinyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl, 1-isindolyl, 3-isindolyl, 4-isindolyl, 5-isindolyl, 6-isindolyl, 7-isindolyl, 2-furyl, 3-furyl, 2-benzofuryl, 3-benzofuryl, 4-benzofuryl, 5-benzofuryl, 6-benzofuryl, 7-benzofuryl, 1-isobenzofuryl, 3-isobenzofuryl, 4-isobenzofuryl, 5-isobenzofuryl, 6-isobenzofuryl, 7-isobenzofuryl, 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl, 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 6-quinoxalyl, 1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl, 9-carbazolyl, 1-phenanthridinyl, 2-phenanthridinyl, 3-phenanthridinyl, 4-phenanthridinyl, 6-phenanthridinyl, 7-phenanthridinyl, 8-phenanthridinyl, 9-phenanthridinyl, 10-phenanthridinyl, 1-acridinyl, 2-acridinyl, 3-acridinyl, 4-acridinyl, 9-acridinyl, 1,7-phenanthroline-2-yl, 1,7-phenanthroline-3-yl, 1,7-phenanthroline-4-yl, 1,7-phenanthroline-5-yl, 1,7-phenanthroline-6-yl, 1,7-phenanthroline-8-yl, 1,7-phenanthroline-9-yl, 1,7-phenanthroline-10-yl, 1,8-phenanthroline-2-yl, 1,8-phenanthroline-3-yl, 1,8-phenanthroline-4-yl, 1,8-phenanthroline-5-yl, 1,8-phenanthroline-6-yl, 1,8-phenanthroline-7-yl, 1,8-phenanthroline-9-yl, 1,8-phenanthroline-10-yl, 1,9-phenanthroline-2-yl, 1,9-phenanthroline-3-yl, 1,9-phen-

anthroline-4-yl, 1,9-phenanthroline-5-yl, 1,9-phenanthroline-6-yl, 1,9-phenanthroline-7-yl, 1,9-phenanthroline-8-yl, 1,9-phenanthroline-10-yl, 1,10-phenanthroline-2-yl, 1,10-phenanthroline-3-yl, 1,10-phenanthroline-4-yl, 1,10-phenanthroline-5-yl, 2,9-phenanthroline-1-yl, 2,9-phenanthroline-3-yl, 2,9-phenanthroline-4-yl, 2,9-phenanthroline-5-yl, 2,9-phenanthroline-6-yl, 2,9-phenanthroline-7-yl, 2,9-phenanthroline-8-yl, 2,9-phenanthroline-10-yl, 2,8-phenanthroline-1-yl, 2,8-phenanthroline-3-yl, 2,8-phenanthroline-4-yl, 2,8-phenanthroline-5-yl, 2,8-phenanthroline-6-yl, 2,8-phenanthroline-7-yl, 2,8-phenanthroline-9-yl, 2,8-phenanthroline-10-yl, 2,7-phenanthroline-1-yl, 2,7-phenanthroline-3-yl, 2,7-phenanthroline-4-yl, 2,7-phenanthroline-5-yl, 2,7-phenanthroline-6-yl, 2,7-phenanthroline-8-yl, 2,7-phenanthroline-9-yl, 2,7-phenanthroline-10-yl, 1-phenazinyl, 2-phenazinyl, 1-phenothiazinyl, 2-phenothiazinyl, 3-phenothiazinyl, 4-phenothiazinyl, 1-phenoxazinyl, 2-phenoxazinyl, 3-phenoxazinyl, 4-phenoxazinyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-oxadiazolyl, 5-oxadiazolyl, 3-furazanyl, 2-thienyl, 3-thienyl, 2-methylpyrrole-1-yl, 2-methylpyrrole-3-yl, 2-methylpyrrole-4-yl, 2-methylpyrrole-5-yl, 3-methylpyrrole-1-yl, 3-methylpyrrole-2-yl, 3-methylpyrrole-4-yl, 3-methylpyrrole-5-yl, 2-t-butylpyrrole-4-yl, 3-(2-phenylpropyl)pyrrole-1-yl, 2-methyl-1-indolyl, 4-methyl-1-indolyl, 2-methyl-3-indolyl, 4-methyl-3-indolyl, 2-t-butyl-1-indolyl, 4-t-butyl-1-indolyl, 2-t-butyl-3-indolyl, or 4-t-butyl-3-indolyl group.

**[0052]** The substituted or unsubstituted alkoxy carbonyl group is represented by -COOY, where Y is, for example, methyl, ethyl, propyl, isopropyl, n-butyl, s-butyl, isobutyl, t-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 2-hydroxyisobutyl, 1,2-dihydroxyethyl, 1,3-dihydroxyisopropyl, 2,3-dihydroxy-t-butyl, 1,2,3-trihydroxypropyl, chloromethyl, 1-chloroethyl, 2-chloroethyl, 2-chloroisobutyl, 1,2-dichloroethyl, 1,3-dichloroisopropyl, 2,3-dichloro-t-butyl, 1,2, 3-trichloropropyl, bromomethyl, 1-bromoethyl, 2-bromoethyl, 2-bromoisobutyl, 1,2-dibromoethyl, 1,3-dibromoisopropyl, 2,3-dibromo-t-butyl, 1,2,3-tribromopropyl, iodomethyl, 1-iodoethyl, 2-iodoethyl, 2-iodoisobutyl, 1,2-diiodoethyl, 1,3-diiodoisopropyl, 2,3-diiodo-t-butyl, 1,2,3-triiodopropyl, aminomethyl, 1-aminoethyl, 2-aminoethyl, 2-aminoisobutyl, 1,2-diaminoethyl, 1,3-diaminoisopropyl, 2,3-diamino-t-butyl, 1,2,3-triaminopropyl, cyanomethyl, 1-cyanoethyl, 2-cyanoethyl, 2-cyanoisobutyl, 1,2-dicyanoethyl, 1,3-dicyanoisopropyl, 2,3-dicyano-t-butyl, 1,2,3-tricyanopropyl, nitromethyl, 1-nitroethyl, 2-nitroethyl, 2-nitroisobutyl, 1,2-dinitroethyl, 1,3-dinitroisopropyl, 2,3-dinitro-t-butyl, or 1,2,3-trinitropropyl group.

**[0053]** The substituted or unsubstituted allylene group is, for example, an aromatic hydrocarbon or condensed polycyclic hydrocarbon such as a substituted or unsubstituted benzen, naphthalene, anthracene, phenanthrene, naphthacene, pyrene, biphenyl or terphenyl; or a divalent group of heterocyclic or condensed heterocyclic compounds such as a substituted or unsubstituted carbazole, pyrrole, thiophene, furan, imidazole, pyrazole, isothiazole, isooxazole, pyridine, pyrazine, pyrimidine, pyridazine, furazane, thianthrene, isobenzofuran, phenoxazine, indolizine, indole, isoindole, 1H-indazole, purine, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole,  $\beta$ -carbazoline, phenanthridine, acridine, perimidine, phenanthroline, phenazine, phenothiazine, or phenoxazine.

**[0054]** The trivalent substituted or unsubstituted aromatic hydrocarbon is a trivalent group of aromatic hydrocarbon or condensed polycyclic hydrocarbon such as a substituted or unsubstituted benzene, naphthalene, anthracene, phenanthrene, naphthacene, pyrene, biphenyl or terphenyl. The trivalent substituted or unsubstituted aromatic heterocyclic group is a trivalent group of heterocyclic or condensed heterocyclic compound such as a substituted or unsubstituted unsubstituted carbazole, pyrrole, thiophene, furan, imidazole, pyrazole, isothiazole, isooxazole, pyridine, pyrazine, pyrimidine, pyridazine, furazane, thianthrene, isobenzofuran, phenoxazine, indolizine, indole, isoindole, 1H-indazole, purine, quinoline, isoquinoline, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole,  $\beta$ -carbazoline, phenanthridine, acridine, perimidine, phenanthroline, phenazine, phenothiazine, or phenoxazine.

**[0055]** The substituted or unsubstituted alkylene group is, for example, a bivalent group of alkane, which may contain one or more substituents, such as a methane, ethane, propane, n-butane, 2-methylpropan, n-pentane, 2-methylbutane, 2,2-dimethylpropane, n-hexane, 2-methyl-n-pentane, 3-methyl-n-pentane, 2,2-dimethylbutane, or 2,3-dimethylbutane. The substituted or unsubstituted alkenylene group is, for example, a bivalent group of alkane, which may contain one or more substituents, such as an ethylene, propylene, 1-butene, 2-butene, or 1,3-butadiene. The substituted or unsubstituted cycloalkylene is, for example, a bivalent group of cycloalkane, which may contain one or more substituents, such as a cyclopropane, cyclobutane, cyclopentane, or cyclohexane.


**[0056]** Examples of the substitutes, which these multivalent groups have, are halogens, hydroxy, substituted or unsubstituted amino, nitro, cyano, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted alkoxy, substituted or unsubstituted aromatic hydrocarbon, substituted or unsubstituted aromatic heterocyclic, substituted or unsubstituted aralkyl, substituted or unsubstituted aryloxy, substituted or unsubstituted alkoxy carbonyl, and carboxy groups. Examples of the bivalent groups, which form a cyclic structure, are tetramethylene, pentamethylene, hexamethylene, diphenylmethane-2,2'-diyl, diphenylmethane-3,3'-diyl, diphenylmethane-4,4'-diyl, and 1,3-butadienyl-1,4-en groups. Metals, which form oxime metal complexes are, for example, aluminium, beryllium, bismuth, cadmium, cerium, cobalt, copper, iron, gallium, germanium, mercury, indium, lanthanum, magnesium, molybdenum, niobium, antimony, scandium, tin, tantalum, thorium, titanium, uranium, tungsten, zirconium, vanadium and zinc. Such host luminescent agents are, for example, the compounds represented by Formulae 62 to 92.

Chemical Formula 62:

5

10

15




Chemical Formula 63:

20

25

30



Chemical Formula 64:


35

40


45

50


55




Chemical Formula 65:



Chemical Formula 66:



Chemical Formula 67:




Chemical Formula 68:

5

10

15




Chemical Formula 69:

20

25

30

35




40

Chemical Formula 70:


45

50


55



Chemical Formula 71:



Chemical Formula 72:



Chemical Formula 73:


5

10

15

20

25



Chemical Formula 74:

30


35

40

45


50

55







Chemical Formula 75:




Chemical Formula 76:




Chemical Formula 77:




Chemical Formula 78:




Chemical Formula 79:




Chemical Formula 80:




Chemical Formula 81:




Chemical Formula 82:




Chemical Formula 83:




Chemical Formula 84:




Chemical Formula 85:




Chemical Formula 86:




Chemical Formula 87:




Chemical Formula 88:




Chemical Formula 89:




Chemical Formula 90:



Chemical Formula 91:




Chemical Formula 92:




[0057] The reference numeral 5 represents an electron injection/ transportation layer which is formed by attaching closely to luminous layer 4 by the same methods as used in anode 2, and then forming a layer having a thickness of 10-500 nm with one or more organic compounds which have a high electron affinity and do not absorb light in a red region, for example, the same compounds as used in luminous layer 4, cyclic ketones such as benzoquinone, anthraquinone, freorenon, or derivatives thereof; silazane derivatives; and conductive oligomers or polymers which have aniline, thiophen, or pyrrole as a repeating unit. In the case of using different materials for electron injection/transportation layers, the materials may be uniformly mixed to form a monolayer or, without mixing, may be separately formed into layers of alternatingly-arranged electron injection/ transportation layers. Examples of respective materials for electron injection/transportation layers are the compounds of Formulae 76, 78, 80, 81, and 93-96.

Chemical Formula 76




Chemical Formula 78:







Chemical Formula 80:




Chemical Formula 81:




Chemical Formula 93:




Chemical Formula 94:



Chemical Formula 95:



Chemical Formula 96:



[0058] The reference numeral 6 represents a cathode which is usually formed by attaching closely to the electron injection/transportation layer 5 and then evaporating thereunto metals such as lithium, magnesium, calcium, sodium, lithium, silver, copper, aluminium, indium, or scandium having a lower work-function usually of, not higher than 6 eV, which is lower than that of the compound used in the electron injection/transportation layer 5; a metal oxide or conductive compound, independently or in combination. The thickness of cathode 6 is not specifically restricted and is set to 10 nm or more, desirably, 50-500 nm, to give a resistivity of 1 k $\Omega/\square$  or lower, in view of electric conductivity, production cost, thickness of devices, and light transmittance. An interface layer, which comprises a compound such as aromatic diamine, quinacridone, naphthalene, organic silicon, or organic phosphide, can be formed to improve the close adhesion between cathode 3 and the electron injection/transportation layer 5, if necessary.

[0059] Thus, the organic EL devices of the present invention can be obtained by forming on substrate 1 uniform layers of anode 2, cathode 6, and luminous layer 4, and optionally, hole injection/transportation layer 3 and electron injection/transportation layer 5 in this order while the adjacent layers are attached together. In forming each layer, it is desirable to consistently operate under a relatively-high vacuum conditions, particularly,  $10^{-5}$  Torr or lower, to suppress the oxidation and decomposition of organic compounds and the adhesion of oxygen and water. To form a luminous layer, the ratio of a host luminous agent and the pyran derivative of the present invention can be regulated by premixing them in a prescribed ratio, or separately controlling their heating velocity in their vacuum evaporation. To suppress the deterioration under use conditions, the organic EL devices thus obtained can be preferably treated in such a manner that a part or the whole of the devices is sealed with a sealing glass or metallic cap under atmosphere of inactivated gas, or covered with a protection layer such as an ultraviolet-hardened resin.

[0060] Explaining the use of the organic EL devices of the present invention, they are driven by intermittently energizing a pulse voltage with a relatively-high voltage, or continuously energizing a non-pulse voltage with a relatively-constant voltage, usually, 3-50 V, dependently on uses. The organic EL devices of the present invention emit light only when the potential of anode exceeds the one of cathode. Thus, the voltage for energizing the organic EL devices of the present invention can be both a direct and alternating currents, and the wave form and the cycle of energizing voltages can be appropriately selected. When energized with an alternating current, the organic EL devices of the present invention alternatively increase and decrease their brightness, or repeatedly turn on and off. In the case of the organic device as shown in FIG.2, when a voltage is energized between anode 2 and cathode 6, holes injected from anode 2 come

into luminous layer 4 through hole injection/transportation layer 3, and electrons injected from cathode 6 come into luminous layer 4 through electron injection/transportation layer 5, respectively. As a result, the holes and electrons rebind in luminous layer 4, then the desired red light is emitted from pyran derivatives in an excited state, generated by the emission through anode 2 and substrate 1. Depending on the kinds of host luminescent agents and the pyran derivatives used in combination, the organic EL devices of the present invention usually have luminous maxima in a red region with wavelengths of 600-670 nm, desirably, 620-660 nm. On an xy-color-degree-table, x of the luminescence is usually in the range of 0.50-0.72 and Y of the luminescence is in the range of 0.20-0.36.

**[0061]** Since the organic EL devices of the present invention have a good color purity of luminescence in a red region and an excellent emission efficiency and durability, they have various uses in luminous bodies and information displaying devices for displaying information visually. Since luminous bodies, which have the organic EL devices of the present invention as a source of light, have a relatively-small consumptive electric power and can be formed into light weight- and flat-boards, they are useful as low-energy and compact-space lighting sources for liquid crystal devices, duplicating machines, inkers, electronic photography machines, computers and the applied machines, industrial controllers, electronic measuring instruments, analyzers, general meters, communications apparatuses, electronic measuring instruments for medical treatment; machines carried on a car, ship, airplane and spaceship; controllers of airplane, interior, signboard, and mark. When used the organic EL devices of the present invention in information displaying devices such as computers, televisions, videos, games, clocks, telephones, car navigations, oscilloscopes, radars, and sonars, they can be driven singly or in combination with organic EL devices, which emit light in a green or blue region, by applying a conventional simple matrix method or active matrix method, if necessary.

**[0062]** When used in dye lasers, the pyran derivatives of the present invention is enclosed in dye cells of laser oscillating machines after purified similarly as in dye-laser active substances, then dissolved in appropriate solvents, and if necessary, followed by adjusting the resulting solutions to an appropriate pH level. As compared with conventional pyran derivatives, the pyran derivatives of the present invention have characteristics that they attain a satisfactory stage gain over a relatively-wide range of wavelengths in visible region, have a relatively-high light-resistance, and do not substantially deteriorate even when used for a relatively-long period of time.


**[0063]** The preferred embodiments according to the present invention are described hereinafter with reference to the following Examples:

#### Example 1


##### Pyran derivative

**[0064]** 1.0 g of the compound represented by Chemical Formula 97 with a 4-dicyanomethylene-2-methyl-4H-pyran skeleton and 1.0 g of the compound represented by Chemical Formula a 98 with a 3-formyl coumarin skeleton were dissolved in five milliliters of N,N-dimethylformamide. The solution was admixed with 0.5 ml piperidine, and the mixture was refluxed for two hours, then cooled to ambient temperature and filtered to collect the formed crystal. The crystal was recrystallized in N, N-dimethylformamide to obtain 0.7 g of a black crystal of the pyran derivative, represented by Chemical Formula 2.

Chemical Formula 97:



Chemical Formula 98:




[0065] Conventional measurement revealed that the pyran derivative had a melting point of 288-290 °C and exhibited an absorption- and fluorescent-maxima at wavelengths of 491 nm and 626 nm, respectively, when measured in methylene chloride. Upon measurement of <sup>1</sup>H-nuclear magnetic resonance spectrum (hereinafter abbreviated as "<sup>1</sup>H-NMR") in chloroform deuteride, the compound showed peaks at chemical shift  $\delta$  (ppm, TMS) of 1.24 (6H, t), 3.06 (6H, s), 3.45 (4H, q), and 6.3-7.7 (14H, m).

### Example 2

#### Pyran derivative

[0066] 1.2 g of the compound represented by Chemical Formula 97 with a 4-dicyanomethylene-2-methyl-4H-pyran skeleton and 2.0 g of the compound represented by Chemical Formula 99 with a 3-formyl coumarin skeleton were dissolved in five milliliters of N, N-dimethylformamide. The solution was admixed with 0.5 ml piperidine, and the mixture was refluxed for two hours, then cooled to ambient temperature and filtered to collect the formed crystal. The crystal was recrystallized in N,N-dimethylformamide to obtain 0.8 g of a black crystal of the pyran derivative, represented by Chemical Formula 5.

## Chemical Formula 99:




**[0067]** Conventional measurement revealed that the pyran derivative had a melting point of 280-282 °C and exhibited an absorption- and fluorescent-maxima at wavelengths of 503 nm and 656 nm, respectively, when measured in methylene chloride. Upon measurement of <sup>1</sup>H-NMR in chloroform deuteride, the compound showed peaks at chemical shift δ (ppm, TMS) of 1.32 (6H, s), 1.57 (6H, s), 1.80 (4H, m), 3.06 (6H, s), 3.34 (4H, m), and 6.4-7.7 (12H, m).

Example 3Pyran derivative

**[0068]** 0.8 g of the compound represented by Chemical Formula 100 with a 4-dicyanomethylene-2-methyl-4H-pyran skeleton and 1.1 g of the compound represented by Chemical Formula 99 with a 3-formyl coumarin skeleton were dissolved in five milliliters of N, N-dimethylformamide. The solution was admixed with 0.5 ml piperidine, and the mixture was refluxed for two hours, then cooled to ambient temperature and filtered to collect the formed crystal. The crystal was recrystallized in N,N-dimethylformamide to obtain 0.4 g of a red-black crystal of the pyran derivative, represented by Chemical Formula 17.

## Chemical Formula 100:




**[0069]** Conventional measurement revealed that the pyran derivative had a melting point of 195-197 °C and exhibited an absorption- and fluorescent-maxima at wavelengths of 509 nm and 662 nm, respectively, when measured in methylene chloride. Upon measurement of <sup>1</sup>H-NMR in chloroform deuteride, the compound showed peaks at chemical shift δ (ppm, TMS) of 0.96 (6H, t), 1.32 (6H, s), 1.58 (6H, s), 1.80 (8H, m), 2.90 (4H, m), 3.4 (4H, m), and 6.3-7.7 (12H, m).

Example 4Pyran derivative

- 5 [0070] 0.8 g of the compound represented by Chemical Formula 101 with a 4-dicyanomethylene-2-methyl-4H-pyran skeleton and 1.2 g of the compound represented by Chemical Formula 98 with a 3-formyl coumarin skeleton were dissolved in seven milliliters of N, N-dimethylformamide. The solution was admixed with 0.7 ml piperidine, and the mixture was refluxed for two hours, then cooled to ambient temperature and filtered to collect the formed crystal. The crystal was recrystallized in N,N-dimethylformamide to obtain 1.0 g of a black crystal of the pyran derivative, represented by
- 10 Chemical Formula 36.

Chemical Formula 101:




- 15
- 20
- 25
- [0071] Conventional measurement revealed that the pyran derivative had a melting point of 329-331 °C and exhibited an absorption- and fluorescent-maxima at wavelengths of 500 nm and 618 nm, respectively, when measured in methylene chloride. Upon measurement of <sup>1</sup>H-NMR in chloroform d<sub>2</sub>, the compound showed peaks at chemical shift δ (ppm, TMS) of 1.24 (12H, t), 3.45 (8H, q), and 6.5-7.7 (14H, m).

Example 5Pyran derivative

- 35
- 40 [0072] 1.0 g of the compound represented by Chemical Formula 102 with a 4-dicyanomethylene-2-methyl-4H-pyran skeleton and 1.3 g of the compound represented by Chemical Formula 99 with a 3-formyl coumarin skeleton were dissolved in four milliliters of N, N-dimethylformamide. The solution was admixed with 0.4 ml piperidine, and the mixture was refluxed for two hours, then cooled to ambient temperature. 0.7 g of the formed crystal was filtered to collect and recrystallized in N,N-dimethylformamide to obtain 0.5 g of a black crystal of the pyran derivative, represented by Chemical Formula 24.

Chemical Formula 102:




[0073] Conventional measurement revealed that the pyran derivative had a melting point of 263-266 °C and exhibited an absorption- and fluorescent-maxima at wavelengths of 519 nm and 677 nm, respectively, when measured in methylene chloride. Upon measurement of <sup>1</sup>H-NMR in chloroform d<sub>2</sub>, the compound showed peaks at chemical shift δ (ppm, TMS) of 1.32 (6H, s), 1.54 (6H, s), 1.7-2.0 (8H, m), 2.8 (4H, m), 3.3 (8H, m), and 6.5-7.7 (10H, m).

#### Example 6

##### Pyran derivative

[0074] 0.6 g of the compound represented by Chemical Formula 103 with a 4-dicyanomethylene-2-methyl-4H-pyran skeleton and 0.5 g of the compound represented by Chemical Formula 99 with a 3-formyl coumarin skeleton were dissolved in three milliliters of N, N-dimethylformamide. The solution was admixed with 0.2 ml piperidine, and the mixture was refluxed for two hours, then cooled to ambient temperature. 1.2 g of the formed crystal was filtered to collect and recrystallized in N,N-dimethylformamide to obtain 1.0 g of a black-green crystal of the pyran derivative, represented by Chemical Formula 53.

Chemical Formula 103:



[0075] Conventional measurement revealed that the pyran derivative had a melting point of 177-180 °C and exhibited an absorption- and fluorescent-maxima at wavelengths of 534 nm and 679 nm, respectively, when measured in methylene chloride. Upon measurement of <sup>1</sup>H-NMR in chloroform d<sub>2</sub>, the compound showed peaks at chemical


shift  $\delta$  (ppm, TMS) of 0.69 (6H, t), 1.32 (6H, s), 1.6 (10H, m), 1.7-1.8 (4H, m), 3.3-3.5 (8H, m), and 6.5-8.5 (16H, m).

### Example 7

#### 5 Pyran derivative

[0076] 1.0 g of the compound represented by Chemical Formula 104 with a 4-dicyanomethylene-2-methyl-4H-pyran skeleton and 0.9 g of the compound represented by Chemical Formula 99 with a 3-formyl coumarin skeleton were dissolved in five milliliters of N, N-dimethylformamide. The solution was admixed with 0.3 ml piperidine, and the mixture  
 10 was refluxed for one hour, then cooled to ambient temperature. 1.6 g of the formed crystal was filtered to collect and recrystallized in N,N-dimethylformamide to obtain 1.4 g of a green-black crystal of the pyran derivative, represented by Chemical Formula 54.

15 Chemical Formula 104:



35 [0077] Conventional measurement revealed that the pyran derivative had a melting point of 249-252 °C and exhibited an absorption- and fluorescent-maxima at wavelengths of 534 nm and 725 nm, respectively, when measured in methylene chloride. Upon measurement of  $^1\text{H-NMR}$  in chloroform  $d_3$ , the compound showed peaks at chemical shift  $\delta$  (ppm, TMS) of 1.32 (6H, s), 1.54 (6H, s), 1.7-2.0 (8H, m), 2.8 (4H, m), 3.3 (8H, m), and 6.3-7.7 (14H, m).

40 [0078] The properties of the pyran derivatives in Examples 1 to 7 are tabulated in Table 1. In Table 1, the wavelengths of absorption maxima were measured after dissolving the pyran derivatives in methylene chloride, and their fluorescent spectra were measured in methylene chloride at a concentration of  $10^{-7}\text{M}$ . As a control, a conventional pyran derivative, represented by Chemical Formula 97, was used.

45

50

55



Table 1

Compound	Maximum absorption wavelength (nm)	Fluorescence		Melting point (°C)	Remarks
		Maximum wavelength (nm)	Relative intensity		
Formula 2	491	626	0.3	288~290	Present invention
Formula 5	503	656	3.0	280~282	Present invention
Formula 17	509	662	1.4	195~197	Present invention
Formula 24	519	677	3.1	263~266	Present invention
Formula 54	534	725	2.2	249~252	Present invention
Formula 53	534	679	0.9	177~180	Present invention
Formula 36	500	618	4.9	329~331	Present invention
Formula 97	468	589	1.8	215~217	Control

[0079] As evident from the results in Table 1, the absorption maxima of the pyran derivatives of the present invention clearly shifted to the side of a longer wavelength as compared with the control, and were close to the oscillating wavelength of argon ion laser and the second harmonic wavelength of YAG laser. Then, the wavelength of absorption

maximum of the pyran derivative, represented by Chemical Formula 2, was close to the oscillating wavelength of 488 nm by argon ion laser; and those of the pyran derivatives, represented by Chemical Formulae 5, 17, 24, 36, 53 and 54, were close to that of the second harmonic wavelength of 532 nm by YAG laser. This demonstrates that the pyran derivatives of the present invention are useful as photosensitizers which polymerize polymerizable compounds with visible light. As far as the fluorescent measurement under the above conditions, fluorescence of the pyran derivatives represented by Chemical Formula 2 was negligible, however, the other pyran derivatives in Table 1 had remarkable luminous abilities and showed equal or significantly-stronger luminescence as compared with those of control. This indicates that the pyran derivatives of the present invention are useful as laser-active substances in dye lasers and luminescent agents in organic EL devices. Although materials and reaction conditions are varied in some degrees dependently on the structures of the pyran derivatives of the present invention, all of the pyran derivatives of the present invention, including the compounds represented by Chemical Formulae 1 to 55, can be easily produced by the methods in Examples 1 and 7 and in accordance therewith.

#### Example 8

##### Photopolymerizable composition

[0080] According to a usual manner, four types of photopolymerizable compositions were prepared by mixing 900 parts by weight of ethyl cellosolve with 100 parts by weight of pentaerythritol acrylate as a photopolymerizable compound, 100 parts by weight of acrylic acid/methacrylic acid copolymer as a binder, eight parts by weight of 3,3', 4,4'-tetra(t-butylperoxycarbonyl) benzophenone as a polymerization initiator, and two parts by weight of any one of pyran derivatives, represented by Chemical Formulae 2, 5, 17 and 36, as a photosensitizer, obtained by the methods in Examples 1 to 4.

[0081] According to a usual manner, the photopolymerizable compositions thus obtained were homogeneously coated over sand-set aluminum plates, which the surfaces had been treated, to form photosensitive layers, followed by forming polyvinyl alcohol layers thereupon to protect the polymerization inhibition by oxygen. A grace scale was closely attached to either of the photosensitive layers and, after installing a 150 W xenon lamp, was irradiated by the lamp with a light of 488 nm, corresponding to the light of argon ion laser, and a light of 532 nm, corresponding to the light of the second harmonic by YAG laser, which were both obtained by combining sharp cut filters ("Y47" and "Y52") commercialized by Toshiba Glass Kabushiki Kaisha, Tokyo, Japan; interference filters ("KL49" and "KL54") commercialized by Toshiba Glass Kabushiki Kaisha, Tokyo, Japan; and a heat-ray-cut-filter ("HA30") commercialized by HOYA Corporation, Tokyo, Japan. Thereafter, the irradiated layers were in a usual manner developed in an alkaline developer, followed by calculating the sensitivity based on the step number at which the compositions were photo-cured in such a manner that, in the numerical formula represented by the following equation, transmittance  $T_n$  of the step number "n" of a step tablet, exposure time t, and exposure strength to were respectively substituted for the equation. Equation:

$$E(\text{mJ}/\text{cm}^2) = \log(\text{mJ}/\text{cm}^2 \text{ s}) \times T_n \times t(\text{s})$$

[0082] In parallel, in place of the pyran derivatives of the present invention, a control system using a conventional compound, represented by Chemical Formula 104, was provided and treated similarly as above. The results are in Table 2.

Table 2

Compound	Measurement at wave-length(nm)	Sensitivity (mJ/cm <sup>2</sup> )	Remarks
Formula 2	488	0.7	Present invention
Formula 5	488	0.6	Present invention
Formula 17	488	0.8	Present invention
Formula 36	488	0.9	Present invention
Formula 97	488	1.4	Control
Formula 2	532	1.5	Present invention
Formula 5	532	1.3	Present invention
Formula 17	532	0.9	Present invention

Table 2 (continued)

Compound	Measurement at wave-length(nm)	Sensitivity (mJ/cm <sup>2</sup> )	Remarks
Formula 36	532	1.4	Present invention
Formula 97	532	2.8	Control

**[0083]** As evident from the results in Table 2, the pyran derivatives of the present invention, represented by Chemical Formulae 2, 5, 17 and 36, showed a higher sensitivity than that of the conventional pyran derivative represented by Chemical Formula 97, when measured at the wavelengths of 488 nm and 532 nm. Particularly, the pyran derivatives of the present invention, represented by Chemical Formulae 2 and 5, had an about two-fold higher sensitivity than that of the conventional pyran derivative represented by Chemical Formula 97, when measured at the wavelength of 488 nm. The pyran derivatives of the present invention, represented by Chemical Formulae 2 and 17, had an about two-fold higher sensitivity than that of the conventional pyran derivative represented by Chemical Formula 97, when measured at the wavelength of 532 nm. The data demonstrate that the pyran derivatives of the present invention are useful to sensitize polymerizable compounds and polymerization initiators in photochemical polymerization.

#### Example 9

##### High-purity pyran derivatives

**[0084]** The pyran derivatives of the present invention, obtained by the methods in Examples 1 to 7, are in a conventional manner purified by sublimation to obtain seven types of high-purity pyran derivatives represented by Chemical Formulae 2, 5, 17, 24, 36, 53, and 54.

**[0085]** The pyran derivatives of this Example are very useful as laser action substances in dye lasers and luminous agents in organic EL devices.

#### Example 10

##### Organic EL device

**[0086]** As shown in FIG. 2, using the pyran derivative of the present invention represented by Chemical Formula 2 as a luminous agent, organic EL devices were prepared by sequentially laying anode 2, hole injection/transportation layer 3, luminous layer 4, electron injection/transportation layer 5, and cathode 6 in this order. Namely, ITO was sputtered on glass substrate 1 to form a membrane as anode 2, having the sheet resistance of 20  $\Omega/\square$ . Anode 2 was evaporated under vacuum conditions with N,N'-diphenyl-N,N'-bis (3-methylphenyl)-[1,1'-biphenyl]-4,4'-diamine represented by Chemical Formula 57 to form a layer, 40 nm in thickness, as hole injection/transportation layer 3, which was closely attached to anode 2; and the compounds represented by Chemical Formulae 76 and 2 were coevaporated on hole injection/transportation layer 3 in a weight ratio of 49 : 1 to form a layer, 50 nm in thickness, as luminous layer 4, which was closely attached to hole injection/transportation layer 3. Further luminous layer 4 was evaporated under vacuum conditions with a compound represented by Chemical Formula 76 to form a layer, 20 nm in thickness, as electron injection/transportation layer 5, which was closely attached to luminous layer 4. Thereafter, on electron injection/transportation layer 5 were co-evaporated magnesium and silver to form a layer, 150 nm in thickness, in an evaporation weight ratio of 10 : 1, as a cathode, which was closely attached to electron injection/transportation layer 5. When a dc current voltage of eight volts was charged between anode 2 and cathode 6 thus obtained, it was emitted a red luminescence having a brightness of 492 cd/m<sup>2</sup> and a luminescent maximum at around a wavelength of 610 nm.

#### Example 11

##### Organic EL device

**[0087]** An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, replacing the pyran derivative represented by Chemical Formula 2 with the one represented by Chemical Formula 5. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 719 cd/m<sup>2</sup> and a luminescent maximum around a wavelength of 620 nm.

Example 12Organic EL device

- 5 [0088] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, replacing the pyran derivative represented by Chemical Formula 26 with the one represented by Chemical Formula 2. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence which had a brightness of 848 cd/m<sup>2</sup> and a luminescent maximum around a wavelength of 623 nm.

10

Example 13Organic EL device

- 15 [0089] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, replacing the pyran derivative represented by Chemical Formula 36 with the one represented by Chemical Formula 2. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 885 cd/m<sup>2</sup>.

20 Example 14Organic EL device

- 25 [0090] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, replacing the pyran derivative represented by Chemical Formula 55 with the one represented by Chemical Formula 2. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 1,010 cd/m<sup>2</sup>.

Example 15

30

Organic EL device

- [0091] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, replacing the pyran derivative represented by Chemical Formula 8 with the one represented by Chemical Formula 2. When energized with a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 622 cd/m<sup>2</sup>.

35

Example 1640 Organic EL device

- [0092] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, using a co-evaporating membrane made of the compound represented by Chemical Formula 58 and the pyran derivative represented by Chemical Formula 36, where the one represented by Chemical Formula 36 was set to a concentration of 5 w/w %. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 867 cd/m<sup>2</sup>.

45

Example 17

50

Organic EL device

- [0093] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, using a co-evaporating membrane made of the compound represented by Chemical Formula 82 and the pyran derivative represented by Chemical Formula 55, where the one represented by Chemical Formula 55 was set to a concentration of 2 w/w %. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 1,218 cd/m<sup>2</sup>.

55

Example 18Organic EL device

5 [0094] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, using a co-evaporating membrane made of the compound represented by Chemical Formula 87 and the pyran derivative represented by Chemical Formula 26, where the one represented by Chemical Formula 26 was set to a concentration of 2 w/w %. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 398

10  $\text{cd/m}^2$ .

Example 19Organic EL device

15 [0095] An organic EL device was prepared similarly as in Example 10 except for, in the preparation of luminous layer 4 in Example 10, using a co-evaporating membrane made of the compound represented by Chemical Formula 90 and the pyran derivative represented by Chemical Formula 55, where the one represented by Chemical Formula 55 was set to a concentration of 0.5 w/w %. When a dc current voltage of eight volts was charged between the anode and the cathode of the organic EL device thus obtained, it was emitted a red luminescence having a brightness of 975

20  $\text{cd/m}^2$ .

Example 20Organic EL device

[0096] As shown in FIG. 2, using the pyran derivative of the present invention represented by Chemical Formula 2 as a luminous agent, organic EL devices were formed by sequentially laying anode 2, hole injection/transportation layer 3, luminous layer 4, electron injection/transportation layer 5, and cathode 6 in this order. Namely, ITO was sputtered on glass substrate 1 to form a membrane, having the sheet resistance of  $20 \Omega/\square$ , as anode 2. Anode 2 was evaporated under vacuum conditions with N,N'-diphenyl-N,N'-bis (1-naphthyl) -[1,1'-biphenyl]-4,4'-diamine represented by Chemical Formula 58 to form a layer, 40 nm in thickness, as hole injection/transportation layer 3, which was closely attached to anode 2. Thereafter, the compounds represented by Chemical Formulae 73 and 2 were co-evaporated on hole injection/transportation layer 3 in a weight ratio of 99 : 1 to form a layer, 50 nm in thickness, as luminous layer 4, which was

30 closely attached to hole injection/transportation layer 3. Further, luminous layer 4 was evaporated under vacuum conditions with the compound represented by Formula 95 to form a layer, 20 nm in thickness, as electron injection/transportation layer 5, which was closely attached to luminous layer 4. Thereafter, electron injection/transportation layer 5 was co-evaporated with magnesium and silver in an evaporation weight ratio of 10 : 1 to form a layer, 150 nm in thickness, as a cathode, which was closely attached to electron injection/transportation layer 5. When a dc current voltage of eight volts was energized between the resulting anode 2 and cathode 6, it was emitted a red luminescence having a brightness of  $451 \text{ cd/m}^2$ .

40

INDUSTRIAL APPLICABILITY

45 [0097] As described above, the present invention was made based on the creation of novel pyran derivatives and the finding of their industrially useful properties. Since the pyran derivatives have absorption maxima in visible region and substantially absorb visible light, they can be very useful as photosensitizers and have various uses such as photochemical polymerization by irradiating visible light such as sun light, carbon arc, high-pressure mercury lamp, xenon lamp, metalhalide lamp, luminescent lamp, tungsten lamp, argon ion laser, krypton ion laser, helium-cadmium laser, helium-neon laser, and the second harmonic wavelength by YAG laser, etc., to photopolymerizing compounds, in the presence or absence of polymerization initiators. Accordingly, the pyran derivatives of the present invention can be very useful in a variety of fields such as plates for printings in laser facsimiles, laser printers, color scanners, black and white scanners, typesettings for photography, plate-makings for flat-board, plate-makings for photogravure, plate-makings for flexo, and plate-makings for silk screen, etc; printings, publications, electrons, information-recordings, chemicals, metals, cars, shipbuildings, and medical treatments, which require photosensitization by visible light as an essential technical element.

55

[0098] Because the pyran derivatives of the present invention have distinctive luminous ability in visible region, particularly, in a red region, they can be very useful as host luminescent agents in organic EL devices, guest luminescent


agents to improve the luminescent efficiency and the luminescent spectrum by doping other host luminescent agents in a negligible amount, and laser-active substances in dye lasers. In particular, the organic EL devices, comprising the pyran derivatives of the present invention, give luminescence with a good color-purity in visible region, particularly, in a red region. This is presumed that the pyran derivatives of the present invention have one or two amino groups in their molecules, and the amino group (s) satisfactorily keeps their amorphous feature in a glass condition and effectively suppresses the cohesion of materials.

[0099] The present invention with such outstanding effects and functions is a significant invention that will greatly contribute to this art.


## 10 Claims

1. A pyran derivative represented by either Formula 1 or 2:

Formula 1:



Formula 2:




where  $R_1$  to  $R_4$  independently represent a straight- or branched-chain of alkyl, alkenyl, or aromatic hydrocarbon group which may have a substituent;  $R_5$  and  $R_6$  independently represent a cyano group, halogenated alkyl group, carboxylic group or substituent derived from carbonic acid, or monocyclic or polycyclic heterocycle group, wherein the heterocycle group may have a substituent;  $R_7$  and  $R_8$  independently represent hydrogen, halogen, alkyl or halogenated alkyl, cyano, or carboxylic group or a substituent derived from carbonic acid;  $R_9$  to  $R_{11}$  independently represent hydrogen or alkyl group;  $R_1$  and  $R_2$ , or  $R_3$  and  $R_4$  may form a cyclic structure along with the nitrogen to which they bound together and with benzene ring bonded to the nitrogen, respectively.

2. The pyran derivative of claim 1, which have an absorption maximum in the visible region.
3. The pyran derivative of claim 1, which have a luminescent maximum in the visible region.
4. A photosensitizer which comprises the pyran derivative of any one of claims 1 to 3.
5. The photosensitizer of claim 4, which sensitizes visible light.
6. A photopolymerizable composition which comprises the pyran derivative of any one of claims 1 to 3.
7. The photopolymerizable composition of claim 6, which comprises a polymerization initiator and/or a binding resin,


and optionally polymerizable compound.

8. The photopolymerizable composition of claim 6 or 7, which can be polymerized by irradiating visible light.
9. A luminescent agent for an organic electro-luminescent device which comprises the pyran derivative of any one of claims 1 to 3.
10. A laser active substance which comprises the pyran derivative of any one of claims 1 to 3.
11. A process for producing a pyran derivative of any one of claims 1 to 3, comprising the step of reacting a compound, represented by Formula 3 having  $R_1$ ,  $R_2$ ,  $R_5$ ,  $R_6$ ,  $R_{10}$  and  $R_{11}$  in Formula 1, with a compound represented by Formula 4 having  $R_3$ ,  $R_4$ ,  $R_7$ , and  $R_9$  in Formula 1:

Formula 3:




Formula 4:




12. A process for producing a pyran derivative of any one of claims 1 to 3, comprising the step of reacting a compound, represented by Formula 5 having  $R_1$ ,  $R_2$ ,  $R_5$ ,  $R_6$ ,  $R_9$  and  $R_{10}$  in Formula 2, with a compound represented by Formula 4 having  $R_3$ ,  $R_4$ ,  $R_7$ , and  $R_9$  in Formula 2:

Formula 4:




Formula 5:




13. A process for producing a pyran derivative of any one of claims 1 to 3, comprising the step of reacting a compound, represented by Formula 6 having R<sub>5</sub> and R<sub>6</sub> in Formula 2, with a compound represented by Formula 4 having R<sub>3</sub>, R<sub>4</sub>, R<sub>7</sub> and R<sub>9</sub> in Formula 2 and a compound represented by Formula 7 having R<sub>1</sub>, R<sub>2</sub>, R<sub>8</sub> and R<sub>10</sub> in Formula 2:




Formula 4:



Formula 6:



Formula 7:



14. An organic electro-luminescent device which comprises the pyran derivative of any one of claims 1 to 3.


15. The organic electro-luminescent device of claim 14, which forms a luminescent layer by using the pyran derivative of any one of claims 1 to 3.

16. The organic electro-luminescent device of claim 14 or 15, wherein the luminescent layer comprises an appropriate host luminescent agent and the pyran derivative of any one of claims 1 to 3.

17. The organic electro-luminescent device of any one of claims 14 to 16, which have a hole injection/transportation layer or an electron injection/transportation layer.

18. The organic electro-luminescent device of any one of claims 14 to 17, which comprises a compound represented by Formula 8 and the pyran derivative of any one of claims 1 to 3.


Formula 8:



where,  $R_{12}$  to  $R_{14}$  independently represent an aromatic hydrocarbon or aromatic heterocyclic group, which may contain a substituent and bind each other to form a cyclic structure.

19. The organic electro-luminescent device of any one of claims 14 to 17, which comprises a compound represented by Formula 9 and the pyran derivative of any one of claims 1 to 3.


Formula 9:



where,  $R_{15}$  to  $R_{18}$  independently represent an aromatic hydrocarbon or aromatic heterocyclic group, which may contain a substituent and bind each other to form a cyclic structure; and Y represents an arylene group which may contain a substituent.

20. The organic electro-luminescent device of any one of claims 14 to 17, which comprises a compound represented by Formula 10 and the pyran derivative of any one of claims 1 to 3.

Formula 10:




where,  $R_{19}$  to  $R_{24}$  independently represent a substituent selected from the group consisting of hydrogen, hydroxy, nitro, cyano, carboxy, amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxycarbonyl groups, where the amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxycarbonyl groups, may contain a substituent;  $L_1$  represents alkyl, alkenyl, cycloalkyl, aromatic hydrocarbon, aromatic heterocyclic, or aralkyl group, where the alkyl, alkenyl, cycloalkyl, aromatic hydrocarbon, aromatic heterocyclic, or aralkyl group may contain a substituent; m represents an integer

selected from 0 to 2, and n represents an integer selected from 1 to 3; and M represents a metal ion with a valency of (n+m).

21. The organic electro-luminescent device of any one of claims 14 to 17, which comprises a compound represented by Formula 11 and the pyran derivative of any one of claims 1 to 3.


Formula 11:



where,  $R_{25}$  to  $R_{36}$  independently represent a substituent selected from the group consisting of hydrogen, halogen, hydroxy, nitro, cyano, carboxy, amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxy carbonyl groups, where the amino, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, aralkyl, aryloxy, and alkoxy carbonyl groups, may contain a substituent;  $L_2$  represents alkylene, alkenylene, cycloalkylene, or a bivalent of aromatic hydrocarbon and aromatic heterocyclic groups, where the alkylene, alkenylene, cycloalkylene, aromatic hydrocarbon, or aromatic heterocyclic group may contain a substituent; 1 represents an integer selected from 0 to 1; s represents an integer selected from 1 to 2; M represents a metal ion with a valency of (s+1).

22. The organic electro-luminescent device of any one of claims 14 to 17, which comprises a compound represented by Formula 12 and the pyran derivative of any one of claims 1 to 3.

Formula 12:



where,  $R_{37}$  to  $R_{40}$  independently represent a substituent selected from the group consisting of cyano, alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, and aryloxy groups, where the alkyl, alkenyl, cycloalkyl, alkoxy, aromatic hydrocarbon, aromatic heterocyclic, and aryloxy groups may contain a substituent and bind each other to form a cyclic structure; Y represents an arylene group which may contain a substituent.

23. The organic electro-luminescent device of any one of claims 14 to 22, which emits a red light.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP00/01437

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IntCl7 C07D407/06, 405/14, 407/14, 417/14, 455/04, 491/16, C09B57/02, C08F2/50, G03C1/73, G03F7/031, G03H1/02		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IntCl7 C07D407/06, 405/14, 407/14, 417/14, 455/04, 491/16, C09B57/02, C08F2/50, G03C1/73, G03F7/031, G03H1/02		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS (STN) REGISTRY (STN)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP, 624580, A1 (Canon Inc.), 17 November, 1994 (17.11.94) & JP, 6-324615, A & US, 5879837, A	1-23
A	EP, 627426, A1 (Canon Inc.), 07 December, 1994 (07.12.94) & JP, 6-329654, A & US, 5492942, A & US, 5723633, A	1-23
A	JP, 8-337582, A (Mitsui Toatsu Chemicals Inc.), 24 December, 1996 (24.12.96) (Family: none)	1-23
A	JP, 10-45741, A (Mitsui Toatsu Chemicals Inc.), 17 February, 1998 (17.02.98) (Family: none)	1-23
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search 07 June, 2000 (07.06.00)		Date of mailing of the international search report 20.06.00
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**